

$(\delta, \chi_{\text{FF}})$ -bounded families of graphs

Manouchehr Zaker*

Department of Mathematics,
Institute for Advanced Studies in Basic Sciences,
Zanjan 45137-66731, Iran

Abstract

For any graph G , the First-Fit (or Grundy) chromatic number of G , denoted by $\chi_{\text{FF}}(G)$, is defined as the maximum number of colors used by the First-Fit (greedy) coloring of the vertices of G . We call a family $\mathcal{F}$ of graphs $(\delta, \chi_{\text{FF}})$ -bounded if there exists a function $f(x)$ with $f(x) \rightarrow \infty$ as $x \rightarrow \infty$ such that for any graph G from the family one has $\chi_{\text{FF}}(G) \geq f(\delta(G))$, where $\delta(G)$ is the minimum degree of G . We first give some results concerning $(\delta, \chi_{\text{FF}})$ -bounded families and obtain a few such families. Then we prove that for any positive integer ℓ , $\text{Forb}(K_{\ell, \ell})$ is $(\delta, \chi_{\text{FF}})$ -bounded, where $K_{\ell, \ell}$ is complete bipartite graph. We conjecture that if G is any C_4 -free graph then $\chi_{\text{FF}}(G) \geq \delta(G) + 1$. We prove the validity of this conjecture for chordal graphs, complement of bipartite graphs and graphs with low minimum degree.

Mathematics Subject Classification (2000): 05C15, 05C07, 05C85, 05C38

Keywords: graph coloring; First-Fit coloring; Grundy number; lower bound; minimum degree

1 Introduction

All graphs in this paper are simple undirected graphs. A family $\mathcal{F}$ of graphs is said to be (δ, χ) -bounded if there exists a function $f(x)$ satisfying $f(x) \rightarrow \infty$ as $x \rightarrow \infty$, such that for any graph G from the family one has $f(\delta(G)) \leq \chi(G)$, where $\delta(G)$ and $\chi(G)$ denotes the minimum degree and chromatic number of G , respectively. Equivalently, the family $\mathcal{F}$ is (δ, χ) -bounded if $\delta(G_n) \rightarrow \infty$ implies $\chi(G_n) \rightarrow \infty$ for any sequence $G_1, G_2, \dots$ with $G_n \in \mathcal{F}$. Motivated by Problem 4.3 in [4], the author introduced and studied (δ, χ) -bounded families of graphs (under the name of δ -bounded families) in [9]. The so-called color-bound family of graphs mentioned in the related problem of [4] is a family for which there exists a function $f(x)$ satisfying $f(x) \rightarrow \infty$ as $x \rightarrow \infty$, such that for any graph G from the family one has $f(\text{col}(G)) \leq \chi(G)$, where $\text{col}(G)$ is defined as $\text{col}(G) = \max\{\delta(H) : H \subseteq G\} + 1$. It was shown in [9] that if we restrict ourselves to hereditary (i.e. closed under taking induced subgraph) families then the two concepts

*mzaker@iasbs.ac.ir

(δ, χ) -bounded and color-bound are equivalent. The first specific results concerning (δ, χ) -bounded families appeared in [9] where the following theorem was proved (in a somewhat different but equivalent form). In the following theorem for any set $\mathcal{C}$ of graphs, $\text{Forb}(\mathcal{C})$ denotes the class of graphs that contains no member of $\mathcal{C}$ as an induced subgraph.

Theorem 1. ([9]) *For any set $\mathcal{C}$ of graphs, $\text{Forb}(\mathcal{C})$ is (δ, χ) -bounded if and only if there exists a constant $c = c(\mathcal{C})$ such that for any bipartite graph $H \in \text{Forb}(\mathcal{C})$ one has $\delta(H) \leq c$.*

Theorem 1 shows that to decide whether $\text{Forb}(\mathcal{C})$ is (δ, χ) -bounded we may restrict ourselves to bipartite graphs. A comprehensive study of (δ, χ) -bounded families was done in [2], where the authors proved the following theorem.

Theorem 2. ([2]) *Given a finite set of graphs $\{H_1, H_2, \dots, H_k\}$. Then $\text{Forb}(H_1, H_2, \dots, H_k)$ is (δ, χ) -bounded if and only if one of the following holds:*

- (i) *For some i , H_i is a star tree.*
- (ii) *For some i , H_i is a forest and for some $j \neq i$, H_j is complete bipartite graph.*

The following result concerns $\text{Forb}(\mathcal{C})$, where $\mathcal{C}$ contains infinitely many graphs in which one of them is a tree.

Theorem 3. ([2]) *Let T be any non star tree. Then $\text{Forb}(T, H_1, \dots)$ is (δ, χ) -bounded if and only if at least one of H_i -s is complete bipartite graph.*

For other (δ, χ) -bounded families of graphs we refer the reader to [2] and [9]. In this paper we work on the Grundy (or First-Fit) chromatic number of graphs. A *Grundy k -coloring* of a graph G , is a proper k -coloring of vertices in G using colors $\{1, 2, \dots, k\}$ such that for any two colors i and j with $i < j$, any vertex colored j is adjacent to some vertex colored i . The Grundy or *First-Fit chromatic number* of a graph G , denoted by $\chi_{\text{FF}}(G)$ (also denoted by $\Gamma(G)$ in some articles), is the largest integer k , such that there exists a Grundy k -coloring for G . It can be shown that $\chi_{\text{FF}}(G)$ is the same as the maximum number of colors used by the First-Fit (greedy) coloring of the vertices of G [7]. To determine $\chi_{\text{FF}}(G)$ is NP-complete even for complement of bipartite graphs G [7]. For this reason it is natural to obtain lower and upper bounds for $\chi_{\text{FF}}(G)$ in terms of ordinary graph theoretical parameters. In this paper we obtain some lower bounds in terms of the minimum degree of graphs. The Grundy number and First-Fit coloring of graphs were studied widely in the literature, see [7, 8] and its references. Throughout the paper we denote the complete graph on n vertices by K_n and the cycle on n vertices by C_n . For each positive integer ℓ , the complete bipartite graph in which each part has ℓ vertices is denoted by $K_{\ell, \ell}$.

2 Results

Generalizing the concept of (δ, χ) -bounded graph, we define the following notion. A family $\mathcal{F}$ of graphs is called $(\delta, \chi_{\text{FF}})$ -bounded if there exists a function $f(x)$ with $f(x) \rightarrow \infty$ as $x \rightarrow \infty$ such that for any graph G from the family one has $\chi_{\text{FF}}(G) \geq f(\delta(G))$. It was shown in [7] that $\chi_{\text{FF}}(G) = 2$ if and only if G is a complete bipartite graph. Obviously, complete bipartite graphs may have arbitrary large minimum degree. We conclude that the family of complete bipartite graphs is not $(\delta, \chi_{\text{FF}})$ -bounded. This example induces that perhaps C_4 and other complete bipartite graphs have significant role in $(\delta, \chi_{\text{FF}})$ -boundedness. Note also that any (δ, χ) -bounded family is also $(\delta, \chi_{\text{FF}})$ -bounded. Another interesting chromatic-related parameter is the so-called coloring number of graphs. The coloring number of a graph G is defined as $\text{col}(G) = \max_{H \subseteq G} \delta(H) + 1$. The coloring number of graphs is a polynomial time parameter. See [4, 5, 9] for more results on the coloring number of graphs. The possible relationships between the coloring number and Grundy number of graphs is an interesting research area. For some graphs G we have $\text{col}(G) < \chi_{\text{FF}}(G)$. For example the path on four vertices P_4 and infinitely many trees satisfy this inequality. Also, for some graphs G we have $\chi_{\text{FF}}(G) < \text{col}(G)$. For example consider complete bipartite graphs $K_{a,b}$, where $a, b \geq 2$. We have the following remark, where by a hereditary family we mean any family $\mathcal{F}$ such that for any graph G from $\mathcal{F}$, if H is an induced subgraph of G then $H \in \mathcal{F}$.

Remark 1. *Let $\mathcal{F}$ be any hereditary family of graphs such that $\mathcal{F}$ is $(\delta, \chi_{\text{FF}})$ -bounded. Then there exists a function $f(x)$ with $f(x) \rightarrow \infty$ as $x \rightarrow \infty$ such that for any graph G from the family one has $\chi_{\text{FF}}(G) \geq f(\text{col}(G))$.*

Proof. Let $G \in \mathcal{F}$ and H be any induced subgraph of G with $\text{col}(G) = \delta(H) + 1$. We have $H \in \mathcal{F}$. Let $g(x)$ be such that $\chi_{\text{FF}}(H) \geq g(\delta(H))$. The proof completes by taking $f(x) = g(x - 1)$. $\square$

As we mentioned before any (δ, χ) -bounded family is also $(\delta, \chi_{\text{FF}})$ -bounded. In Theorem 5 we obtain $(\delta, \chi_{\text{FF}})$ -bounded families which are not (δ, χ) -bounded. For this purpose we first obtain in Proposition 1 a result concerning (δ, χ) -boundedness of graphs. In the following, the girth of a graph G is the length of a shortest cycle contained in G . When G contains a cycle then we say that G has finite girth. We use also the following two facts. The first fact states that any graph with m edges contains a bipartite subgraph with at least $m/2$ edges. The second one states that any graph with n vertices and $m/2$ edges contains a subgraph with minimum degree at least $m/2n$. For the proof of these facts we refer the reader to standard Graph Theory books such as [1].

Proposition 1. *Let $\mathcal{C}$ be any finite collection of graphs such that any member of $\mathcal{C}$ has finite girth. Then $\text{Forb}(\mathcal{C})$ is not (δ, χ) -bounded. In particular $\text{Forb}(K_3, K_{2,m})$ and $\text{Forb}(K_{\ell, \ell})$ are not (δ, χ) -bounded.*

Proof. Let g be an even integer such that the girth of any graph in $\mathcal{C}$ is at most g . For the proof we use the following Turán-type result which is attributed to Erdős in [6]. For

any k and n there exists a graph on n vertices with $\Omega(n^{1+1/2k-1})$ edges that contains no cycle of length at most $2k$. Let $g = 2k$, and recall from the previous paragraph that (1) a graph with m edges contains a bipartite subgraph with at least $m/2$ edges, and (2) a graph with n vertices and $m/2$ edges contains a subgraph with minimum degree at least $m/2n$. We conclude that there exists an infinite sequence $G_1, G_2, \dots$ of bipartite graphs such that $\delta(G_i) \rightarrow \infty$ as $i \rightarrow \infty$ and the girth of any G_i is more than g . This shows that G_i belongs to $\text{Forb}(\mathcal{C})$. This shows that $\text{Forb}(\mathcal{C})$ is not (δ, χ) -bounded. $\square$

In opposite direction we show in Theorem 5 that $\text{Forb}(K_3, K_{2,m})$ is $(\delta, \chi_{\text{FF}})$ -bounded. More generally, Theorem 7 asserts that $\text{Forb}(K_{\ell, \ell})$ is $(\delta, \chi_{\text{FF}})$ -bounded. Before we proceed, we need to introduce a family of trees T_k , $k = 1, 2, \dots$. For $k = 1, 2$, T_1 (resp. T_2) is isomorphic to K_1 (resp. K_2). Assume that T_k has been defined. Attach a leaf to any vertex of T_k and denote the resulting tree by T_{k+1} . It is easily observed that $\chi_{\text{FF}}(T_k) = k$. Note also that $|V(T_k)| = 2^{k-1}$. We need also the following result from ([9], Theorem 2), where $\rho(G) = |E(G)|/|V(G)|$.

Theorem 4. ([9]) *Let G be any triangle-free graph such that G does not contain $K_{2,m}$, where $m > 1$. If $\rho(G) \geq (k-3)(m-1) + 1$ then G contains all trees on k vertices as induced subgraphs.*

The promised result is as follows.

Theorem 5. *Let $G \in \text{Forb}(K_3, K_{2,m})$. Then*

$$\chi_{\text{FF}}(G) \geq \log\left(\frac{\delta(G) + 6m - 8}{2m - 2}\right) + 1.$$

Proof. First note that G does not contain triangle and $K_{2,m}$. Set $\delta(G) = p$ and $k = (p + 6m - 8)/(2m - 2)$, for simplicity. We have

$$\begin{aligned} (k-3)(m-1) + 1 &= \left(\frac{p + 6m - 8}{2m - 2} - 3\right)(m-1) + 1 \\ &= \left(\frac{p - 2}{2m - 2}\right)(m-1) + 1 \\ &= \frac{p}{2}. \end{aligned}$$

Since $\rho(G) \geq (p/2)$ then G satisfies the conditions of Theorem 5 with these k and m . Therefore G contains all trees on k vertices as induced subgraph. In particular G contains T_q as induced subgraph, where $q = \log((\delta(G) + 6m - 8)/(2m - 2)) + 1$. We conclude that

$$\chi_{\text{FF}}(G) \geq \log\left(\frac{\delta(G) + 6m - 8}{2m - 2}\right) + 1.$$

$\square$

By applying Theorem 4 and Theorem 5 more economically when $m = 2$ we obtain the following bound.

Corollary 1. *Let $G \in \text{Forb}(K_3, C_4)$. Then*

$$\chi_{\text{FF}}(G) \geq \log(\delta(G) + 1).$$

We noted before that the family of complete bipartite graphs is not $(\delta, \chi_{\text{FF}})$ -bounded. Hence the following proposition is immediate from this fact.

Proposition 2. *Let $\mathcal{C}$ be any collection of graphs such that any member of it contains an odd cycle. Then $\text{Forb}(\mathcal{C})$ is not $(\delta, \chi_{\text{FF}})$ -bounded. In particular $\text{Forb}(K_3)$ is not $(\delta, \chi_{\text{FF}})$ -bounded.*

In Theorem 7 we prove that $\text{Forb}(K_{\ell, \ell})$ is $(\delta, \chi_{\text{FF}})$ -bounded. For this purpose we need the following theorem from [2].

Theorem 6. ([2]) *For every tree T and for positive integers ℓ, k there exist a function $f(T, \ell, k)$ with the following property. If G is a graph with $\delta(G) \geq f(T, \ell, k)$ and $\chi(G) \leq k$ then G contains either T or $K_{\ell, \ell}$ as an induced subgraph.*

We shall make use of this theorem in proving the next result.

Theorem 7. *For each positive integer ℓ , $\text{Forb}(K_{\ell, \ell})$ is $(\delta, \chi_{\text{FF}})$ -bounded.*

Proof. Recall that for each positive integer k , T_k denotes the only smallest tree of Grundy number k . Let $\{G_n\}_{n=1}^{\infty}$ be a sequence of $K_{\ell, \ell}$ -free graphs such that $\delta(G_n) \rightarrow \infty$ as $n \rightarrow \infty$. Assume on the contrary that for some integer N , $\chi_{\text{FF}}(G_n) \leq N$ holds for all n . It follows that for each n , T_{N+1} is not an induced subgraph of G_n . Hence G_n belongs to $\text{Forb}(T_{N+1}, K_{\ell, \ell})$. Theorem 6 implies that for each n either $\delta(G_n) < f(T_{N+1}, \ell, N)$ or $\chi(G_n) > N$. But the second case is impossible because $\chi(G_n) \leq \chi_{\text{FF}}(G_n) \leq N$. Therefore for each n , $\delta(G_n) < f(T_{N+1}, \ell, N)$. This contradicts $\delta(G_n) \rightarrow \infty$. This contradiction completes the proof. $\square$

The following result shows that chordal graphs are $(\delta, \chi_{\text{FF}})$ -bounded with $f(x) = x + 1$. Note that the class of chordal graphs is the same as $\text{Forb}(C_4, C_5, \dots)$. In a graph G , by a simplicial vertex we mean any vertex v such that $G[N(v)]$ is a clique in G , where $G[N(v)]$ stands for the subgraph of G induced by the set $N(v)$ of the neighbors of v in G . It is a known fact that any chordal graph G admits a simplicial elimination ordering (see e.g. [1]). In other words, let G be a chordal graph. Then there exists a vertex ordering $v_1, \dots, v_n$ of G such that v_i is simplicial in $G \setminus \{v_1, \dots, v_{i-1}\}$. We shall make use of this fact in the following theorem.

Theorem 8. *$\text{Forb}(C_4, C_5, \dots)$ is $(\delta, \chi_{\text{FF}})$ -bounded with $f(x) = x + 1$.*

Proof. Let G be any chordal graph G and let $v_1, v_2, \dots, v_n$ be a simplicial ordering of the vertices of G . Since $G[N(v) \cup \{v_1\}]$ is a clique in G with $\deg_G(v_1) + 1$ vertices, then $\omega(G) \geq \deg_G(v) + 1 \geq \delta(G) + 1$. We have also $\chi_{\text{FF}}(G) \geq \chi(G) \geq \omega(G)$. Hence $\chi_{\text{FF}}(G) \geq \delta(G) + 1$. $\square$

By strengthening Theorem 8 we propose the following conjecture.

Conjecture 1. *Let G be a C_4 -free graph. Then $\chi_{\text{FF}}(G) \geq \delta(G) + 1$.*

A natural scenario to prove the above conjecture is as follows. Let $\mathcal{F}$ be a hereditary family of graphs satisfying the following property. Any member G from the family contains a maximal independent set (MIS) such as I such that $\delta(G \setminus I) = \delta(G) - 1$. We have the following observation which can be proved by induction.

Observation 1. *Let $\mathcal{F}$ be any hereditary family of graphs such that any graph G from the family contains a MIS, say I such that for any vertex v of G if $\deg_G(v) = \delta(G)$ then $\deg_{G \setminus I}(v) = \deg_G(v) - 1$. Then $\chi_{\text{FF}}(G) \geq \delta(G) + 1$ for any graph G from $\mathcal{F}$.*

Unfortunately the family of C_4 -free graphs does not satisfy the above condition. In this regard it is worthy to work on the following problem.

Problem. *Find families $\mathcal{F}$ of graphs satisfying the following property. Any graph G from $\mathcal{F}$ contains a MIS, say I such that for any vertex v of G if $\deg_G(v) = \delta(G)$ then $\deg_{G \setminus I}(v) = \deg_G(v) - 1$.*

In Theorem 9 we show that Conjecture 1 holds for any graph which is the complement of a bipartite graph. We need some prerequisites. In a graph H , a subset D of edges in H is called an edge dominating set if each edge in $E(H) \setminus D$ has a common end point with an edge in D . Let H be any bipartite graph. Set $G = \overline{H}$. Let $\gamma'(H)$ be the smallest size of an edge dominating set in H . It was proved in [7] that $\chi_{\text{FF}}(G) = |V(G)| - \gamma'(H)$. We have now the following theorem.

Theorem 9. *Let H be any bipartite graph and G be the complement of H such that G is C_4 -free. Then $\chi_{\text{FF}}(G) \geq \delta(G) + 1$.*

Proof. Let n be the order of G . Since $\delta(G) = n - \Delta(H) - 1$, then the inequality $\chi_{\text{FF}}(G) \geq \delta(G) + 1$ is equivalent to $\gamma'(H) \leq \Delta(H)$. Now we use the fact that for any edge dominating set R in a bipartite graph, there is a matching M which is also an edge dominating set and $|M| \leq |R|$. This fact can be easily proved and we omit mentioning its proof here, and refer the reader to [3]. Let R be an edge dominating set in H with $|R| = \gamma'(H)$. Using the previous fact we obtain that R is a matching and therefore $\gamma'(H) \leq \alpha'(H)$. Hence to complete the proof we need to show that $\alpha'(H) \leq \Delta(H)$. We prove the latter inequality by induction on the number of edges. Note that since G

is C_4 -free then H is $2K_2$ -free, where by $2K_2$ we mean the graph consisting only of two independent edges.

Let M be any matching of maximum size in H and $e = uv$ be any edge of M . Define another graph as $H_0 = H \setminus \{u, v\}$. We have $\alpha'(H_0) \leq \Delta(H_0)$. Hence $\alpha'(H) - 1 \leq \Delta(H_0)$. To finalize the proof we show that $\Delta(H_0) + 1 \leq \Delta(H)$. Otherwise, since H_0 is an induced subgraph of H , we have $\Delta(H_0) = \Delta(H)$. Let x be any vertex in H_0 such that $\deg_{H_0}(x) = \Delta(H)$. Without loss of generality, we may assume that u and x are in the same bipartite part of H . We show that u is adjacent to any neighbor of x . Let w be any neighbor of x . Since H is $2K_2$ -free then the subgraph of H consisting of two edges uv and xw can not be induced in H . Now, since x has the maximum degree then x can not be adjacent to v in H . Hence u should be adjacent to w . But v is adjacent to u and not adjacent to x . This means that the degree of u is strictly greater than the degree of x , a contradiction with our choice of x . This completes the proof. $\square$

The following theorem shows that Conjecture 1 holds for all graphs G with $\delta(G) \leq 3$.

Theorem 10. *Let G be a C_4 -free graph with $\delta(G) \leq 3$. Then $\chi_{\text{FF}}(G) \geq \delta(G) + 1$.*

Proof. Theorem obviously holds if $\delta(G) = 1$. Assume that $\delta(G) = 2$. Let v be any vertex of G and a, b any two neighbors of v . If a and b are adjacent then the resulting triangle shows that $\chi_{\text{FF}}(G) \geq 3$. Assume that a and b are not adjacent. Let c be any neighbor of b . If a and c are not adjacent then we obtain an induced P_4 on the vertex set $\{v, a, b, c\}$. Hence the desired inequality holds in this case. Assume that a and c are adjacent. Then since G is C_4 -free and b is not adjacent to a then v is adjacent to c . This gives rise to a triangle. Hence in this case too $\chi_{\text{FF}}(G) \geq 3$.

Assume now that $\delta(G) = 3$. Let v be any vertex of degree 3. Let a, b, c be the neighbors of v .

Case 1. Three vertices a, b, c are independent.

In this case we first note that no two vertices from $\{a, b, c\}$ have a common neighbor other than the vertex v , since otherwise let u be a common neighbor of a and b . The two vertices a and b are independent and G is C_4 -free. Hence v should be adjacent to u . This contradicts $\deg_G(v) = 3$. Now, let x and y (resp. z and t) be two neighbors of a (resp. b). We have $\{x, y\} \cap \{z, t\} = \emptyset$ and c is not adjacent to any vertex in $\{x, y, z, t\}$. Consider a small bipartite graph H consisting of the bipartite sets $\{x, y\}$ and $\{z, t\}$ with all edges from G among these parts. If there are at most two edges in H then we color v by 4, a by 3, b by 2 and c by 1. The vertex a needs two neighbors of colors 1 and 2; and the vertex b needs one neighbor of color 1. We can easily fulfil these conditions by assigning suitable colors 1 and 2 to the vertices of H . If there are exactly three edges in H then (assuming that x is adjacent to z) we consider the following coloring. We color x by 4, z by 3, y and t by 2 and a and b by 1. Finally, we consider the case that H is a complete bipartite graph. In this case we color x by 4, z by 3, a and b by 2; and t and v by 1 (note that t and v are not adjacent). All of these pre-colorings are partial Grundy colorings with 4 colors. This completes the proof in Case 1.

Case 2. a and b are adjacent and c is not adjacent to a and also to b .

In this case we color v by 4, a by 2, b by 3 and c by 1. Let d be a neighbor of b . We color d by 1. Note that c and d can not be adjacent. If a is adjacent to d then we obtain a partial Grundy coloring using four colors. Otherwise, let a be adjacent to a vertex say e . If e is adjacent to d then since a is not adjacent to d , hence b should be adjacent to e . In this situation we color e by 1 and remove the color of d . Now the colors of $\{v, a, b, c, e\}$ is a partial Grundy coloring with four colors. But if e is not adjacent to d , we color both vertices e and d by 1. Note that in this case the colors of $\{v, a, b, c, d, e\}$ introduce a partial Grundy coloring using four colors.

Case 3. a is adjacent to both b and c ; but b and c are not adjacent.

In this case we color v by 4, a by 3, b by 1 and c by 2. Let d be a new neighbor of c . If b and d are not adjacent then we color both of them by 1. The resulting coloring is a partial Grundy coloring using four colors. But if b and d are adjacent, then a should be adjacent to d . Now consider the 4-cycle on $\{v, b, c, d\}$. Since the degree of v is three then v can not be adjacent to d . Hence b and c should be adjacent. But this is a contradiction.

Case 4. The only remaining case is that a, b, c are all adjacent. But in this case we obtain a clique of size four. It is clear that in this case $\chi_{\text{FF}}(G) \geq 4$. $\square$

We end the paper by mentioning that Conjecture 1 is also valid for graphs with minimum degree four. The proof is by checking too many cases. We omit the details.

References

- [1] R. Diestel, Graph Theory, Springer-Verlag, New York, 2000.
- [2] A. Gyárfás, M. Zaker, On (δ, χ) -bounded graphs, Electron. J. Combin. 18 (2011) #P108.
- [3] R. Haggkvist, A.S. Asratian, T.M.G. Denley, Bipartite Graphs and their Applications, Cambridge University Press, Cambridge, 1998.
- [4] T.R Jensen, B. Toft, Graph Coloring Problems, Wiley, New York, 1995.
- [5] S.E. Markossian, G.S. Gasparian, B.A. Reed, β -perfect graphs, J. Combin. Theory Ser. B 67 (1996) 1-11.
- [6] M. Simonovits, Extremal graph theory, In: Selected Topics in Graph Theory vol. 2 (1983) L. Beineke and R. Wilson Eds., Academic Press: pp. 161–200.
- [7] M. Zaker, Results on the Grundy chromatic number of graphs, Discrete Math. 306 (2006) 3166–3173.
- [8] M. Zaker, Inequalities for the Grundy chromatic number of graphs, Disc. Appl. Math. 155 (2007) 2567–2572.
- [9] M. Zaker, On lower bounds for the chromatic number in terms of vertex degree, Discrete Math. 311 (2011) 1365–1370.